

**¡Feliz cumpleaños, Kathleen Ollerenshaw!**

Escrito por Marta Macho Stadler  
Lunes 01 de Octubre de 2012 09:00

---



**Kathleen Ollerenshaw** es una matemática y política británica, conocida -en el ámbito matemático- por sus importantes trabajos sobre [cuadrados mágicos](#).

Hoy 1 de octubre cumple **100** años, así que esta entrada es un pequeño homenaje a esta mujer, sin duda de una gran fortaleza.


Nació en Manchester (Inglaterra) con grandes problemas de audición y estaba completamente sorda cuando ingresó a estudiar en la [Oxford University](#). Tras doctorarse en matemáticas, se casó con el médico militar Robert Ollerenshaw. Su carrera política comenzó en [Rusholme](#),

, y en la década de 1980, fue asesora de confianza de la Primera Ministra británica [Margaret Thatcher](#).

**¡Feliz cumpleaños, Kathleen Ollerenshaw!**

Escrito por Marta Macho Stadler  
Lunes 01 de Octubre de 2012 09:00

---



The Institute of Mathematics and its Applications

[http://www.maths.org.uk/research/magic\\_squares.html](http://www.maths.org.uk/research/magic_squares.html) El número más grande conocido es el número de Erdős, igual a 5, a través de Hermann Bondi, Ivor



## THEOREM OF THE DAY

**The Ollerenshaw–Brée Formula** Let  $n$ , a positive integer, have prime factorisation  $n = p_1^{s_1} p_2^{s_2} \cdots p_r^{s_r}$  in which  $p_1 = 2$  and  $s_1 \geq 2$ , so that  $n$  is doubly even. The number of  $n \times n$  most-perfect magic squares, up to horizontal, vertical and diagonal symmetry, is given by

$$N(n) = M(n) \sum_{v=0}^{\tau(n)} W(v)[W(v) + W(v+1)],$$

where  $\tau(n)$  is the number of divisors of  $n$  and  $M(n)$  and  $W(v)$  are given by  $M(n) = 2^{n-2}(n/2)!^2$ , and

$$W(v) = \sum_{i=0}^v (-1)^{v+i} \binom{v+1}{i+1} \prod_{j=1}^r \binom{5j+i}{i}.$$

A most-perfect 8×8 square

Suppose the integers  $0, \dots, n^2 - 1$  are arranged in an  $n \times n$  array,  $n$  a multiple of 4. The result is called *most-perfect* if (a) any two entries at distance  $n/2$  on any diagonal sum to  $n^2 - 1$ ; and (b) the entries in any  $2 \times 2$  block of adjacent cells sums to  $2(n^2 - 1)$ . In the  $8 \times 8$  example here, the cells diagonally opposite on the yellow bands (sloping down to the right) are pairwise at distance 4 and sum, pairwise, to 63. The  $2 \times 2$  blocks, including ‘broken’ blocks, like the one which counts the top-left cell as adjacent to bottom-left and top-right and so on, have cells summing to 126. Most-perfect squares are *pandiagonally magic*: each complete row, column and (broken) diagonal (such as the yellow and green diagonals) sum to the same quantity:  $n(n^2 - 1)/2$ . The  $4 \times 4$  example shows how a most-perfect square may be constructed from any *reversible square*, that is, one in which all  $2 \times 2$  submatrices have equal diagonal sums, and in which, in any row or column, the sums (cell  $i$ ) + (cell  $n - i + 1$ ) are equal for all  $i$ ,  $1 \leq i \leq n/2$ . Conversely, every most-perfect square may be obtained in this way.

Kathleen Ollerenshaw was in her 80s when she made the amazing discovery that a 1938 construction of J. Barkley Rosser (of the Church-Rosser property) and Robert J. Walker could be adapted to give precisely the members of this major class of magic squares. Working with David Brée she exploited this to give the first ever counting formula for *any* such class (the total number of all  $n \times n$  magic squares is unknown even for  $n = 6$ ).

An example of order 4

0	1	4	5
2	3	6	7
8	9	12	13
10	11	14	15

0	1	5	4
2	3	7	6
8	9	13	12
10	11	15	14

0	62	2	60	7	57	5	59
15	49	13	51	8	54	10	52
16	46	18	44	23	41	21	43
31	33	29	35	24	38	26	36
56	6	58	4	63	1	61	3
55	9	53	11	48	14	50	12
40	22	42	20	47	17	45	19
39	25	37	27	32	30	34	28

Reverse column order on RHS

0	1	5	4
2	3	7	6
8	9	13	12
10	11	15	14

Reverse row order in bottom half

0	14
7	9
10	4
13	3

Distr 2x2

Web link: [http://recmath.org/Magic\\_Squares/most-perfect.htm](http://recmath.org/Magic_Squares/most-perfect.htm)

Further reading: *Most Perfect Pandiagonal Magic Squares*, by Kathleen Ollerenshaw and David Brée, Institute of Mathematics and its Applications,

From [www.theoremoftheday.org](http://www.theoremoftheday.org) by Robin Whitty. This file hosted by London South Bank University

Digitized by the Internet Archive with funding from the Internet Archive Books Project and the Internet Archive Library Development Fund and the Internet Archive Book Images Project. The Internet Archive is a non-profit organization building a digital library of historical publications that is freely available for everyone to use, enjoy, and share. We believe that everyone should have access to all of the world's knowledge. Since our inception in 1990, we have been a leader in digitizing historical texts and in making these resources free and open to the public. We also serve as a hub for the Internet Archive community, which includes many partners and supporters, like the Internet Archive Foundation, the Internet Archive Library, the Internet Archive Book Images Project, and the Internet Archive Book Images Project. We are grateful for their support and cooperation in helping to build a better future for everyone.